



King's Research Portal

DOI:

[10.1111/jcpp.12958](https://doi.org/10.1111/jcpp.12958)

[Link to publication record in King's Research Portal](#)

Citation for published version (APA):

Hedman, A., Breithaupt, L., Hübel, C., Thornton, L. M., Tillander, A., Norring, C., Birgegård, A., Larsson, H., Ludvigsson, J. F., Säwendahl, L., Almqvist, C., & Bulik, C. M. (2018). Bidirectional relationship between eating disorders and autoimmune diseases. *Journal of child psychology and psychiatry*.
<https://doi.org/10.1111/jcpp.12958>

Citing this paper

Please note that where the full-text provided on King's Research Portal is the Author Accepted Manuscript or Post-Print version this may differ from the final Published version. If citing, it is advised that you check and use the publisher's definitive version for pagination, volume/issue, and date of publication details. And where the final published version is provided on the Research Portal, if citing you are again advised to check the publisher's website for any subsequent corrections.

General rights

Copyright and moral rights for the publications made accessible in the Research Portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognize and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the Research Portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the Research Portal

Take down policy

If you believe that this document breaches copyright please contact librarypure@kcl.ac.uk providing details, and we will remove access to the work immediately and investigate your claim.

Bidirectional relationship between eating disorders and autoimmune diseases

Anna Hedman, Ph.D.¹, Lauren Breithaupt, M.A.^{1,2}, Christopher Hübel, M.D., M.Sc.,^{1,3} Laura M. Thornton, Ph.D.⁴, Annika Tillander, Ph.D.^{1,5}, Claes Norring, Ph.D.⁶, Andreas Birgegård, Ph.D.⁶, Henrik Larsson, Ph.D.^{1,7}, Jonas Ludvigsson, M.D., Ph.D.^{1,8}, Lars Sävendahl, M.D., Ph.D.⁹, Catarina Almqvist, M.D., Ph.D.^{1,10}, Cynthia M. Bulik, Ph.D.^{1,4,11}

¹Department of Medical Epidemiology and Biostatistics, Karolinska Institutet, Stockholm, Sweden

²Department of Psychology, George Mason University, Fairfax, VA, USA

³Social, Genetic & Developmental Psychiatry Centre, Institute of Psychiatry, Psychology & Neuroscience, King's College London, UK

⁴Department of Psychiatry, University of North Carolina at Chapel Hill, Chapel Hill, NC

⁵Department of Computer and Information Science, Linköping University, Linköping, Sweden

⁶Centre for Psychiatry Research, Department of Clinical Neuroscience, Karolinska Institutet, & Stockholm Health Care Services, Stockholm County Council, Stockholm, Sweden

⁷School of Medical Sciences, Örebro University, Örebro, Sweden

⁸Department of Pediatrics, Örebro University Hospital, Örebro, Sweden

⁹Department of Women's and Children's Health, Karolinska Institutet, Stockholm, Sweden

¹⁰Pediatric Allergy and Pulmonology Unit, Astrid Lindgren Children's Hospital, Karolinska University Hospital, Stockholm, Sweden

¹¹Department of Nutrition, University of North Carolina at Chapel Hill, Chapel Hill, NC

Total word count: 5843

Running Title: Associations between eating disorder and autoimmune diseases.

Abstract

Background: Immune system dysfunction may be associated with eating disorders, and associations may have implications for detection, risk assessment, and treatment of both autoimmune diseases and eating disorders. However, the nature of the relationship between these two disease entities remains unclear. We evaluated the strength of associations for the bidirectional relationships between eating disorders and autoimmune diseases. **Methods:** In this nationwide population-based cohort study, Swedish registers were linked to establish a cohort of over two and a half million individuals born in Sweden between January 1, 1979 and December 31, 2005 and followed-up until December 2013. Two Cox proportional hazard regression models were used to investigate 1) subsequent risk of eating disorders in individuals with autoimmune diseases and 2) subsequent risk of autoimmune diseases in individuals with eating disorders. **Results:** We observed a strong, bidirectional relationship between the two classes of illness indicating that diagnosis of one illness increased the risk of the other. In women, autoimmune disease exposure increased subsequent hazard of anorexia nervosa, bulimia nervosa, and other eating disorders. Likewise, anorexia nervosa, bulimia nervosa, and other eating disorders increased subsequent hazard of autoimmune diseases. The gastrointestinal-related autoimmune diseases celiac and Crohn's showed a bidirectional relationship with anorexia nervosa and other eating disorders. Psoriasis showed a bidirectional relationship only with other eating disorders. Prior type 1 diabetes increased risk for anorexia nervosa, bulimia nervosa, and other eating disorders. In men, we did not observe a bidirectional pattern, but prior autoimmune arthritis increased risk for other eating disorders. **Conclusions:** The associations between eating disorders and autoimmune diseases provide support for previously reported associations and the bidirectional risk pattern observed suggests either a shared underlying mechanism or a third explanatory variable contributing to the association of these illnesses.

Keywords: hazard, risk, immune system, cox regression, anorexia nervosa, bulimia nervosa, autoimmunity

Introduction

Autoimmunity has been implicated in several psychiatric disorders (Barbuti et al., 2017; Eaton et al., 2006; Edmiston, Ashwood, & Van de Water, 2017; Frick & Pittenger, 2016; Mechawar & Savitz, 2016; Muskens, Velders, & Staal, 2017; Neigh & Ali, 2016; Tylee et al., 2017; Volk, 2017; see eTable 1 for a review), including eating disorders (Raevuori et al., 2014; Wotton, James, & Goldacre, 2016; Zerwas et al., 2017). Moreover, the first genome-wide significant association in anorexia nervosa (AN) was identified (Duncan et al., 2017) in a region previously implicated in autoimmune diseases, including type 1 diabetes (Barrett et al., 2009) and arthritis (Okada et al., 2014).

Eating disorders and autoimmunity are complex traits influenced by numerous genetic variants acting additively in combination with environmental factors to influence phenotypic expression (Yilmaz, Hardaway, & Bulik, 2015). Autoimmunity varies on a continuum, ranging from no clinical consequences to pathogenic autoimmunity causing inflammatory organ infiltration, tissue damage, and overt disease-specific symptomatology. Autoimmune diseases occur in ~7-9% of the population and increase with age (Theofilopoulos, Kono, & Baccala, 2017).

Extensive Swedish health registers afforded an exploration of bidirectional associations between eating disorders and autoimmune diseases (eTable 2). We extended prior studies (Raevuori et al., 2014; Wotton et al., 2016; Zerwas et al., 2017) by examining a cohort of more than 2.5 million individuals.

Methods

Study Population and Data Sources

We studied individuals aged 0-35 years born in Sweden between January 1, 1979 and December 31, 2005, excluding those who emigrated or died before age 8, or were from multiparous births, to reduce nesting. Individuals were followed until eating disorder onset, autoimmune disease onset, death, emigration from Sweden, or the end of the follow-up period (December 31, 2013), whichever came first. We linked registers using the national personal identification number (Ludvigsson, Otterblad-Olausson, Pettersson, & Ekblom, 2009). Consistent with prior research (Mårild et al., 2017; Mustelin et al., 2017; Yao et al., 2016), birth year and sex were from the Total Population Register (Ludvigsson et al., 2016); migration data were from the Migration Register (Statistics Sweden); causes of death were from the Cause of Death Register (Statistics Sweden); and socioeconomic status (SES) was estimated using highest parental education (i.e., completed year 9 or below; completed year 12; >2 years tertiary) from the Longitudinal Integration Database for Health Insurance and Labour Market Studies (Statistics Sweden, 2012) when the child was 8 years old. The study was approved by the Regional Ethical Review Board in Stockholm, Sweden.

Eating Disorder Outcomes

Eating disorder diagnoses were from the National Patient Register (NPR), which tracks all inpatient care since 1987 and outpatient care since 2001 (Ludvigsson et al., 2011; Center for Epidemiology at the National Board of Health and Welfare, n.d.); Riksät-National Quality Register for Treatment for Eating Disorders (since 1999) (Emilsson, Lindahl, Köster, Lambe, & Ludvigsson, 2015); the regional quality assurance system for eating disorders, Stepwise (since 2005; Birgegård, Björck, & Clinton, 2010); and the clinical database for child and adolescent psychiatry in Stockholm, Pastill (since 2001; Lindevall, 2009). NPR discharge diagnoses were from Swedish ICD-9 and ICD-10 (Smedby, 2006). In Riksät and Stepwise, eating disorder

diagnoses were based on *DSM-IV-TR* (American Psychiatric Association, 2005) (Birgegård et al., 2010). Coverage in Riksät and Stepwise has increased over time (Javaras et al., 2015). Pastill eating disorder diagnoses were based on ICD-10 or DSM-IV from Child and Adolescent Mental Health Services in Stockholm County (Lindevall, 2009).

Four eating disorder outcomes were evaluated: 1) AN: 307B (Swedish ICD-9); F50.0 or F50.1 (ICD-10); or DSM-IV AN or atypical AN; 2) Other eating disorders: 307F (Swedish ICD-9); F50.2, F50.3, or F50.9 (ICD-10); or DSM-IV bulimia nervosa (BN), atypical BN, binge-eating disorder (BED), or eating disorder not otherwise specified (EDNOS). 3) Any eating disorder: included all individuals with an AN, BN, and/or other eating disorder diagnosis. Consistent with previous reports (Allen, Byrne, Oddy, & Crosby, 2013; Stice, Marti, & Rohde, 2013), many individuals had both AN and other eating disorders (at different times) and are included as incident cases for both the AN and other eating disorder outcomes. Thus, the number of incident cases of AN and/or other eating disorders is greater than the number of incident cases of any eating disorder, where each individual can be an incident case only once. Analysis of any eating disorder provide information on the overall incidence of eating disorders without inflation due to diagnostic crossover. 4) BN: included all individuals who received a BN diagnosis in the NPR (F50.2, F50.3) since 1997 or in Riksät, Stepwise, or Pastill. Analyses of BN are considered secondary because the years of observation are fewer than for the other diagnostic groups. BED and EDNOS could not be separated because Swedish ICD-9 only included a heterogeneous category (eating disorders other than AN) and F50.8, from ICD-10, was not consistently used to code BED prior to DSM-5.

Age of onset reflects first contact with the respective diagnosis in the NPR, Riksät, Stepwise, or Pastill after the 8th birthday. The minimum age of onset for eating disorders was 8 to avoid diagnostic misclassification (e.g., childhood feeding difficulties; N excluded=4,493).

Autoimmune Disease Outcome

Autoimmune disease diagnoses were obtained from the NPR using Swedish ICD revisions 8-10 diagnoses based on year of diagnosis. We evaluated any autoimmune disease as a group, and by specific categories: celiac disease, Crohn's disease, ulcerative colitis, psoriasis, arthritis, lupus, and type 1 diabetes (eTable 2). Age of onset was age at first contact with the respective diagnosis in the NPR with no minimum age of onset.

Statistical Analyses

Data management was conducted with SAS version 9.4; analyses were conducted with STATA version 14. False discovery rate (FDR) corrections were made for each predefined set of hypothesis tests (n=128) (Benjamini & Hochberg, 2000; Glickman, Rao, & Schultz, 2014).

To optimize our longitudinal data, we performed two sets of Cox proportional hazard regression models estimating the relative hazards of eating disorders following autoimmune disease exposure and autoimmune disease risk following eating disorder exposure. In the first set, autoimmune disease status (presence/absence) was treated as time-dependent exposure and calendar year (1987-1996, 1997-2006, 2007-2013) was adjusted for as a time varying variable; all other variables were considered time-independent. For each analysis, we estimated hazard ratios (HR) for eating disorders in individuals exposed to an autoimmune disease compared to those not exposed. In the second set of analyses, we estimated HRs for autoimmune diseases, comparing individuals exposed to an eating disorder to those not exposed, with eating disorder status as a time-varying exposure variable. All estimates were adjusted for calendar-time in the parametric part of the Cox model, and for age in the nonparametric part. SES was entered as a covariate into all models. In Cox models, HRs greater than 1 indicate a higher risk of illness, whereas HRs less than 1 indicate a lower risk of illness compared with unaffected individuals. Analyses were applied to males and females separately.

Significantly increased HRs, after FDR correction, for an autoimmune disease after eating disorder exposure were investigated for effects of temporal proximity by calculating HR for ≤ 1 year, >1 year to ≤ 4 years, and >4 years between exposure and outcome. Due to possible misdiagnosis of eating disorders before age 8, we did not explore temporal proximity between autoimmune disease exposure and eating disorder onset. Furthermore, we did not investigate different ages of eating disorder onset as age of onset distributions revealed no evidence of bi- or multimodal patterns justifying such an approach, rendering any age at onset cutoff arbitrary (Figures 1a and 1b).

Results

Our cohort comprised 2,545,611 individuals (51.4% males, 48.6% females) followed over 33,640,644 person-years (range: 1 month to 22 years).

Sample characteristics

Table 1 presents prevalence and age at first diagnosis. Eating disorders occurred more commonly in females (2.0%) than in males (0.1%): 94% of eating disorder cases were female.

The most common autoimmune diseases were celiac disease, type 1 diabetes, and psoriasis. A higher percentage of females than males were diagnosed with an autoimmune disease.

Risk of Eating Disorders Following Exposure to Autoimmune Disease

Tables 2a and 2b provide the results from the Cox regression models estimating HRs of eating disorders after an autoimmune disease exposure in males and females, respectively. The following sections only present significant results.

In males, any preceding autoimmune disease was associated with an 82% increased hazard in other eating disorders, a 78% increased hazard in any eating disorder, and a 56% increased hazard for BN. In females, any preceding autoimmune disease increased the hazard for AN by 59%, for other eating disorders by 71%, for any eating disorder by 62%, and for BN by 57%.

We also evaluated the risk for subsequent eating disorders following exposure to celiac disease, Crohn's disease, ulcerative colitis, psoriasis, arthritis, lupus, and type 1 diabetes. In males, exposure to arthritis increased the risk for other eating disorders by 357% and any eating disorder by 267%, however, the assumption of proportional hazards was violated. In females, the risk for subsequent AN was increased after celiac disease (50%), Crohn's disease (89%), and type 1 diabetes (71%). Moreover, celiac disease (47%), Crohn's disease (63%), ulcerative colitis (52%), psoriasis (33%), and type 1 diabetes (153%) increased risk for subsequent other eating disorders. Risk for any eating disorder was increased after exposure to celiac disease (45%), Crohn's disease (61%), ulcerative colitis (49%), psoriasis (27%), and type 1 diabetes (119%). Risk for subsequent BN was increased by 222% by an earlier diagnosis of type 1 diabetes.

Risk of Autoimmune Disease Following Exposure to an Eating Disorder

Tables 3a and 3b provide the results from the Cox regression models estimating risk for subsequent autoimmune disease following exposure to an eating disorder in males and females. After correcting for multiple testing, we did not observe an increased risk for autoimmune diseases after eating disorder exposure in men. However, females with AN (42%), other eating disorders (58%), any eating disorder (53%), or BN (48%) were at increased risk for later autoimmune diseases.

We evaluated the risk for celiac disease, Crohn's disease, ulcerative colitis, psoriasis, arthritis, lupus, and type 1 diabetes following a diagnosis of AN, other eating disorders, any

eating disorder, or BN in males and females. Males diagnosed with eating disorders did not show an increased risk for subsequent autoimmune diseases after correcting for multiple testing. In females, prior exposure to AN (83%), other eating disorders (69%), and any eating disorder (72%) increased risk for celiac disease. Other eating disorders (72%) and any eating disorder exposure (63%) showed increased risk for Crohn's disease. Exposure to other eating disorders increased the risk for subsequent psoriasis by 38%.

Temporal Proximity of Risk of Autoimmune Disease Following Exposure to an Eating Disorder

We explored effects of temporal proximity between exposure to eating disorders and risk for autoimmune disease in females (eTable 3) as only females had significant findings. Exposure to AN was associated with an increased risk (105%) of being diagnosed with any autoimmune disease within the first year after AN diagnosis and with a 49% increased risk between years 1 and 4. Similarly, risk for celiac disease was increased 217% within the first year and 84% in years 1 to 4 after AN diagnosis. The exposure to other eating disorders increased the risk of being diagnosed with any autoimmune disease for all three time periods: 216% for 1 year after diagnosis; 47% for 1 to 4 years between diagnoses; and 45% for >4 years. Other eating disorders increased the risk of being diagnosed with celiac disease (165%) and Crohn's disease (188%) within the first year. The increased risk for celiac disease persisted between 1 year and 4 years at 55%. Within the first year, females with any eating disorder were at 114% increased risk of developing any autoimmune disease; 48% increased risk between years 1 and 4; and 32% increased risk >4 years. Any eating disorder increased the risk for celiac disease and Crohn's disease within the first year after diagnoses by 189% and 202%, respectively. The risk of being diagnosed with celiac disease persisted between years 1 and 4 at 58%. BN was associated with a 79% increased risk for any autoimmune disease after 4 years of BN diagnosis.

Discussion

With 2.5 million participants and over 26,000 individuals diagnosed with an eating disorder, to our knowledge, this is the largest prospective register-based study examining the bidirectional associations between autoimmune diseases and eating disorders. Consistent with previous research, we observed positive and strong associations between eating disorders and autoimmune diseases that are on par with reported associations in epidemiological investigations between autoimmune diseases and other psychiatric disorders, such as attention-deficit/hyperactivity disorder (Nielsen, Benros, & Dalsgaard, 2017), depression (Andersson et al., 2015; Euesden, Danese, Lewis, & Maughan, 2017), and schizophrenia (Benros et al., 2014; Chen et al., 2012). We extended previous observations by investigating a variety of autoimmune diseases previously associated with eating disorders in case reports, clinical samples, and smaller cohort studies (Wotton et al., 2016). Our results replicate bidirectional risk patterns of eating disorders and autoimmune diseases explored in two clinical cohort studies (Raevuori et al., 2014; Wotton et al., 2016) and a Danish national study (Zerwas et al., 2017).

In men, we did not observe any bidirectional patterns. Preceding autoimmune arthritis and type 1 diabetes increased the risk for other eating disorders and preceding autoimmune arthritis increased the risk for any eating disorder. The UK study also observed a significantly increased risk for BN in males with type 1 diabetes based on reports from five cases (Wotton et al., 2016).

In women, AN showed a bidirectional relationship with celiac disease, replicating reported results (Mårild et al., 2017; Wotton et al., 2016). Crohn's disease increased risk for AN (Raevuori et al., 2014; Wotton et al., 2016), but was not bidirectionally associated (Wotton et al., 2016). The gastrointestinal-related autoimmune diseases celiac and Crohn's showed a bidirectional relationship with other eating disorders. The Finnish study (Raevuori et al., 2014)

showed increased odds for Crohn's disease and BN, whereas the UK study (Wotton et al., 2016) found an increased risk for celiac disease after BN. Ulcerative colitis increased risk for other eating disorders. Additionally, psoriasis showed a bidirectional relationship with other eating disorders, replicating findings from the UK study (Wotton et al., 2016) and clinical case reports (Basavaraj, Navya, & Rashmi, 2011; Crosta et al., 2014; Ferreira, Abreu, Reis, & Figueiredo, 2016). Type 1 diabetes increased risk for AN, other eating disorders, and BN as previously reported (Raevuori et al., 2014; Wotton et al., 2016); however, Wotton et al. (2016) reported an increased risk for type 1 diabetes after AN and BN which we did not replicate.

The bidirectional nature of some associations suggests either a shared underlying mechanism or a third explanatory variable that influences risk for both disease groups. Such risk-elevating factors could be genetic, environmental, or a combination of both. Current evidence suggests that dysregulated immune function may be one shared underlying mechanism. Several biological factors influencing immune function are described to be predominant or altered in eating disorders: a female preponderance (Chowen, Argente-Arizón, Freire-Regatillo, & Argente, 2017; Klein & Flanagan, 2016; McCarthy, Nugent, & Lenz, 2017), metabolic changes mediated by adipokines such as leptin and adiponectin (Abella et al., 2017; de Candia et al., 2017; Nikolajczyk, Jagannathan-Bogdan, Shin, & Gyurko, 2011; Saucillo, Gerriets, Sheng, Rathmell, & Maciver, 2014), elevated cytokines (Solmi et al., 2015), abnormal levels of estrogen (Khan & Ansar Ahmed, 2015; Klump, Culbert, & Sisk, 2017), and lower abundance or diversity of intestinal microbiota (Carr, Kleiman, Bulik, Bulik-Sullivan, & Carroll, 2016; Kleiman et al., 2015; Lam, Maguire, Palacios, & Caterson, 2017; Postler & Ghosh, 2017; Rooks & Garrett, 2016). These factors potentially influence the relationship between eating disorders and autoimmunity. For example, cortisol levels are dysregulated in eating disorders (Monteleone et al., 1999) possibly due to an altered stress response. Cortisol is often included in the therapeutic

regimen for autoimmune diseases (Ilzarbe et al., 2017; Straub & Cutolo, 2016). Furthermore, Fetissov et al. detected autoantibodies against appetite-regulating peptides, including α -Melanocyte-stimulating hormone and adrenocorticotrophic hormone in AN and BN (Fetissov et al., 2002, 2005, 2008). However, the role of autoantibodies in autoimmune diseases is not fully understood. Antibodies may be an epiphenomenon or have a causal effect facilitating aberrant immune cell function leading to cytotoxicity.

Recent research suggests a genetic overlap between several autoimmune diseases and psychiatric disorders; however, the only study to date that has included eating disorders revealed no significant genetic associations between AN and autoimmune diseases or traits (Tylee et al., 2017). The increased risk for eating disorders after type 1 diabetes could be metabolically mediated through a dysregulation of insulin homeostasis, administration of mandatory external insulin, and insulin misuse (Bryden et al., 1999; Colton et al., 2015; Colton, Olmsted, Daneman, Rydall, & Rodin, 2004). Similarly, increased risk of any eating disorder after ulcerative colitis could be due to symptom- and therapy-related behavioral changes as treatment often includes dietary changes and a colectomy with pouches or stoma (Ungaro, Mehandru, Allen, Peyrin-Biroulet, & Colombel, 2017). Patients suffering from inflammatory bowel disease often report eating behavior changes related to their disease, and many perceive food as a risk factor for relapse decreasing their pleasure in eating (Zallot et al., 2013). Moreover, the increased risk of being diagnosed with a gastrointestinal-related autoimmune disease within the first year of having been diagnosed with an eating disorder may suggest diagnostic uncertainty due to the overlap in their clinical presentation complicating the differential diagnosis (Golden & Park, 2017; Ilzarbe et al., 2017; Tokatly Latzer et al., 2018).

Environmental factors, such as diet, dietary behavior, and smoking, also influence the human immune system. Starvation and food restriction (as seen in AN) could reduce

inflammation and attenuate symptoms in autoimmune illnesses (Hafström, Ringertz, Gyllenhammar, Palmblad, & Harms-Ringdahl, 1988). For gastrointestinal-associated immune diseases, dietary changes are often prescribed to control pain, diarrhea, and bleeding (Lane, Zisman, & Suskind, 2017). All of these processes may be active, and the inclusion of genetic, biological, and environmental confounders, as well as a developmental perspective in prospective studies are needed to further clarify the relationships between eating disorders and autoimmune diseases.

Strengths and Limitations

Study strengths include the total-population design and substantially more eating disorder cases than in previous studies (Mårild et al., 2017; Raevuori et al., 2014; Wotton et al., 2016; Zerwas et al., 2017). Prospectively collected data allowed us to explore bidirectional risk and minimized the risk of recall bias since data were routinely collected blind to the hypothesis of this study. For most chronic diseases in the inpatient register, the positive predictive value was found to differ between diagnoses in the NPR and medical journals, but is generally 85-95% (Ludvigsson et al., 2011). In addition, we investigated specific autoimmune diseases, which can inform disease detection, treatment, and management.

The observed bidirectional relationships contribute to evidence of immune system involvement in some subtypes of eating disorders. However, we cannot entirely rule out misdiagnosis or surveillance bias given symptom overlap, especially with gastrointestinal autoimmune diseases. Despite the large sample size and follow-up period, some autoimmune disease risk periods remain outside study timeline (e.g., rheumatoid arthritis age of onset is typically after age 44; Symmons, 2002) and exact ages of onset for diseases cannot be traced in register data. Moreover, although the study covers the peak age of onset for eating disorders

(Javaras et al., 2015), individuals remain at risk across the lifespan. A longer follow-up period could alter some of the bidirectional relationships. Additionally, we were unable to evaluate the associations of BED because it could not be distinguished from other eating disorders in the NPR. The small number of male eating disorder cases limited our investigation of sex differences. Given the large sample size, small differences in prevalence between the sexes would be statistically significant, but their clinical significance questionable. Furthermore, the assumptions of proportional hazards were violated in some of the associations (Tables 2-3). Lastly, the Swedish versions of ICD-8 and ICD-9 could not distinguish between type 1 diabetes and type 2 diabetes; however, as our cohort is young (≤ 34 years), most individuals with ICD-8 or ICD-9 diabetes are assumed to have type 1 diabetes. Diabetes diagnoses according to ICD-10, the largest proportion in our cohort, are classified into type 1 or type 2.

Conclusions

Results clarify the bidirectional relation between eating disorders and autoimmune processes. Clinically, our results encourage vigilance for the emergence of these illnesses in afflicted individuals. As the size of genomic investigations of both eating disorders and autoimmune diseases increases, we will be well positioned to further explore the extent to which shared genetic factors influence risk for both classes of illness. Identifying common environmental risk factors although less tractable, may also be facilitated by the identification of risk variants or profiles.

Key Points

- In the largest prospective register-based study to date on eating disorders and autoimmune diseases, we replicated strong bidirectional relationships between eating disorders and various autoimmune diseases.
- The observed positive and strong associations between eating disorders and

Acknowledgements

This research was supported by the Swedish Research Council (PI: Bulik; VR Dnr: 538-2013-8864); the Anorexia Nervosa Genetics Initiative (ANGI), an initiative of the Klarman Family Foundation; the Foundation of Hope: Research and Treatment of Mental Illness; and the National Science Foundation Graduate Research Fellowship under Grant No.1000183151.

Financial support for the data linkage was provided from the Swedish Research Council through the *Swedish Initiative for Research on Microdata in the Social And Medical Sciences* (SIMSAM) framework grant no 340-2013-5867 and grants provided by the Stockholm County Council (ALF-projects). Any opinion, findings, and conclusions or recommendations expressed in this material are those of the authors(s) and do not necessarily reflect the views of the National Science Foundation.

H. Larsson has served as a speaker for Eli-Lilly and Shire and has received research grants from Shire; all outside the submitted work. Dr. Bulik is a grant recipient from and has served on advisory boards for Shire. She receives royalties from Pearson and Walker. All interests unrelated to this work. Dr. Norring is a consultant on a research grant from Shire, unrelated to this work. L. Sävendahl has served as a speaker for Merck, Novo Nordisk and Pfizer and has received research grant from Merck; all outside the submitted work. L. Breithaupt, C. Almqvist, Dr. Birgegård, Dr. Hedman, Dr. Ludvigsson, Dr. Thornton, Dr. Tillander, and Dr. Hübel have nothing to disclose.

Correspondence to: Dr. Bulik, Department of Psychiatry, University of North Carolina at Chapel Hill, CB #7160, 101 Manning Drive, Chapel Hill, NC 27599-7160, USA; tel: +46-8-524 800 00, e-mail: cbulik@med.unc.edu

References

- Abella, V., Scotece, M., Conde, J., Pino, J., Gonzalez-Gay, M. A., Gómez-Reino, J. Gualillo, O. (2017). Leptin in the interplay of inflammation, metabolism and immune system disorders. *Nature Reviews Rheumatology*, 13, 100–109.
- Allen, K. L., Byrne, S. M., Oddy, W. H., & Crosby, R. D. (2013). DSM--IV--TR and DSM-5 eating disorders in adolescents: Prevalence, stability, and psychosocial correlates in a population-based sample of male and female adolescents. *Journal of Abnormal Psychology*, 122, 720.
- American Psychiatric Association. (2005). *Diagnostic and Statistical Manual of Mental Disorders, Fourth Edition, Text Revision*. American Psychiatric Publishing.
- Andersson, N. W., Gustafsson, L. N., Okkels, N., Taha, F., Cole, S. W., Munk-Jørgensen, P., & Goodwin, R. D. (2015). Depression and the risk of autoimmune disease: A nationally representative, prospective longitudinal study. *Psychological Medicine*, 45, 3559–3569.
- Barbuti, M., Carvalho, A. F., Köhler, C. A., Murru, A., Verdolini, N., Guiso, G., ... Pacchiarotti, I. (2017). Thyroid autoimmunity in bipolar disorder: A systematic review. *Journal of Affective Disorders*, 221, 97–106.
- Barrett, J. C., Clayton, D. G., Concannon, P., Akolkar, B., Cooper, J. D., Erlich, H. A., ... Type 1 Diabetes Genetics Consortium. (2009). Genome-wide association study and meta-analysis find that over 40 loci affect risk of type 1 diabetes. *Nature Genetics*, 41, 703–707.
- Basavaraj, K. H., Navya, M. A., & Rashmi, R. (2011). Stress and quality of life in psoriasis: An update. *International Journal of Dermatology*, 50, 783–792.
- Benjamini, Y., & Hochberg, Y. (2000). On the adaptive control of the false discovery rate in multiple testing with independent statistics. *Journal of Educational and Behavioral Statistics*, 25, 60–83.
- Benros, M. E., Pedersen, M. G., Rasmussen, H., Eaton, W. W., Nordentoft, M., & Mortensen, P. B. (2014). A nationwide study on the risk of autoimmune diseases in individuals with a personal or a family history of schizophrenia and related psychosis. *American Journal of Psychiatry*, 171, 218–226.

- Birgegård, A., Björck, C., & Clinton, D. (2010). Quality assurance of specialised treatment of eating disorders using large-scale Internet-based collection systems: Methods, results and lessons learned from designing the Stepwise database. *European Eating Disorders Review*, 18, 251–259.
- Bryden, K. S., Neil, A., Mayou, R. A., Peveler, R. C., Fairburn, C. G., & Dunger, D. B. (1999). Eating habits, body weight, and insulin misuse. A longitudinal study of teenagers and young adults with type 1 diabetes. *Diabetes Care*, 22, 1956–1960.
- Carr, J., Kleiman, S. C., Bulik, C. M., Bulik-Sullivan, E. C., & Carroll, I. M. (2016). Can attention to the intestinal microbiota improve understanding and treatment of anorexia nervosa? *Expert Review of Gastroenterology & Hepatology*, 10, 565–569.
- Center for Epidemiology at the National Board of Health and Welfare, NBHW. (n.d.) The National Patient Register. Retrieved from <http://www.socialstyrelsen.se/register/halsodataregister/patientregistret/inenglish>
- Chen, S.-J., Chao, Y.-L., Chen, C.-Y., Chang, C.-M., Wu, E. C.-H., Wu, C.-S., ... Tsai, H.-J. (2012). Prevalence of autoimmune diseases in in-patients with schizophrenia: Nationwide population-based study. *British Journal of Psychiatry*, 200, 374–380.
- Chowen, J. A., Argente-Arizón, P., Freire-Regatillo, A., & Argente, J. (2017). Sex differences in the neuroendocrine control of metabolism and the implication of astrocytes. *Frontiers in Neuroendocrinology*, 48, 3–12.
- Colton, P. A., Olmsted, M. P., Daneman, D., Farquhar, J. C., Wong, H., Muskat, S., & Rodin, G. M. (2015). Eating disorders in girls and women with type 1 diabetes: A longitudinal study of prevalence, onset, remission, and recurrence. *Diabetes Care*, 38, 1212–1217.
- Colton, P., Olmsted, M., Daneman, D., Rydall, A., & Rodin, G. (2004). Disturbed eating behavior and eating disorders in preteen and early teenage girls with type 1 diabetes: A case-controlled study. *Diabetes Care*, 27, 1654–1659.
- Crosta, M. L., Caldarola, G., Fraietta, S., Craba, A., Benedetti, C., Coco, V., ... De Simone, C. (2014). Psychopathology and eating disorders in patients with psoriasis. *Giornale Italiano Di Dermatologia E Venereologia*, 149, 355–361.

- de Candia, P., De Rosa, V., Gigantino, V., Botti, G., Ceriello, A., & Matarese, G. (2017). Immunometabolism of human autoimmune diseases: From metabolites to extracellular vesicles. *FEBS Letters*, 591, 3119–3134.
- Duncan, L., Yilmaz, Z., Gaspar, H., Walters, R., Goldstein, J., Anttila, V., ... Bulik, C. M. (2017). Significant locus and metabolic genetic correlations revealed in genome-wide association study of anorexia nervosa. *American Journal of Psychiatry*, 174, 850–858.
- Eaton, W. W., Byrne, M., Ewald, H., Mors, O., Chen, C.-Y., Agerbo, E., & Mortensen, P. B. (2006). Association of schizophrenia and autoimmune diseases: Linkage of Danish national registers. *American Journal of Psychiatry*, 163, 521–528.
- Edmiston, E., Ashwood, P., & van de Water, J. (2017). Autoimmunity, autoantibodies, and autism spectrum disorder. *Biological Psychiatry*, 81, 383–390.
- Emilsson, L., Lindahl, B., Köster, M., Lambe, M., & Ludvigsson, J. F. (2015). Review of 103 Swedish healthcare quality registries. *Journal of Internal Medicine*, 277, 94–136.
- Euesden, J., Danese, A., Lewis, C. M., & Maughan, B. (2017). A bidirectional relationship between depression and the autoimmune disorders - New perspectives from the National Child Development Study. *PloS One*, 12, e0173015.
- Ferreira, B. I. R. C., Abreu, J. L. P. D. C., Reis, J. P. G. D., & Figueiredo, A. M. D. C. (2016). Psoriasis and associated psychiatric disorders: A systematic review on etiopathogenesis and clinical correlation. *Journal of Clinical and Aesthetic Dermatology*, 9, 36–43.
- Fetissov, S. O., Hallman, J., Orelund, L., af Klinteberg, B., Grenbäck, E., Hulting, A.-L., & Hökfelt, T. (2002). Autoantibodies against α -MSH, ACTH, and LHRH in anorexia and bulimia nervosa patients. *Proceedings of the National Academy of Sciences*, 99, 17155–17160.
- Fetissov, S. O., Hamze Sinno, M., Coquerel, Q., Do Rego, J. C., Coëffier, M., Gilbert, D., ... Déchelotte, P. (2008). Emerging role of autoantibodies against appetite-regulating neuropeptides in eating disorders. *Nutrition*, 24, 854–859.

- Fetissov, S. O., Harro, J., Jaanisk, M., Järv, A., Podar, I., Allik, J., ... Hökfelt, T. (2005). Autoantibodies against neuropeptides are associated with psychological traits in eating disorders. *Proceedings of the National Academy of Sciences of the United States of America*, 102, 14865–14870.
- Frick, L., & Pittenger, C. (2016). Microglial dysregulation in OCD, Tourette syndrome, and PANDAS. *Journal of Immunology Research*, 2016, 8606057.
- Glickman, M. E., Rao, S. R., & Schultz, M. R. (2014). False discovery rate control is a recommended alternative to Bonferroni-type adjustments in health studies. *Journal of Clinical Epidemiology*, 67, 850–857.
- Golden, N. H., & Park, K. T. (2017). Celiac disease and anorexia nervosa—an association well worth considering. *Pediatrics*, 139, e20170545.
- Hafström, I., Ringertz, B., Gyllenhammar, H., Palmblad, J., & Harms-Ringdahl, M. (1988). Effects of fasting on disease activity, neutrophil function, fatty acid composition, and leukotriene biosynthesis in patients with rheumatoid arthritis. *Arthritis and Rheumatism*, 31, 585–592.
- Ilzarbe, L., Fàbrega, M., Quintero, R., Bastidas, A., Pintor, L., García-Campayo, J., ... Ilzarbe, D. (2017). Inflammatory bowel disease and eating disorders: A systematized review of comorbidity. *Journal of Psychosomatic Research*, 102, 47–53.
- Javaras, K. N., Runfol, C. D., Thornton, L. M., Agerbo, E., Birgegård, A., Norring, C., ... Bulik, C. M. (2015). Sex- and age-specific incidence of healthcare-register-recorded eating disorders in the complete Swedish 1979-2001 birth cohort. *International Journal of Eating Disorders*, 48, 1070–1081.
- Khan, D., & Ansar Ahmed, S. (2015). The immune system is a natural target for estrogen action: opposing effects of estrogen in two prototypical autoimmune diseases. *Frontiers in Immunology*, 6, 635.
- Kleiman, S. C., Watson, H. J., Bulik-Sullivan, E. C., Huh, E. Y., Tarantino, L. M., Bulik, C. M., & Carroll, I. M. (2015). The intestinal microbiota in acute anorexia nervosa and during renourishment: Relationship to depression, anxiety, and eating disorder psychopathology. *Psychosomatic Medicine*, 77, 969–981.

- Klein, S. L., & Flanagan, K. L. (2016). Sex differences in immune responses. *Nature Reviews Immunology*, 16, 626–638.
- Klump, K. L., Culbert, K. M., & Sisk, C. L. (2017). Sex differences in binge eating: Gonadal hormone effects across development. *Annual Review of Clinical Psychology*, 13, 183–207.
- Lam, Y. Y., Maguire, S., Palacios, T., & Caterson, I. D. (2017). Are the gut bacteria telling us to eat or not to eat? Reviewing the role of gut microbiota in the etiology, disease progression and treatment of eating disorders. *Nutrients*, 9.
- Lane, E. R., Zisman, T. L., & Suskind, D. L. (2017). The microbiota in inflammatory bowel disease: current and therapeutic insights. *Journal of Inflammation Research*, 10, 63–73.
- Lindevall, O. (2009). Pastill—a comprehensive clinical database for child and adolescent psychiatry in Stockholm. *European Society for Child and Adolescent Psychiatry*.
- Ludvigsson, J. F., Almqvist, C., Bonamy, A.-K. E., Ljung, R., Michaëlsson, K., Neovius, M., ... Ye, W. (2016). Registers of the Swedish total population and their use in medical research. *European Journal of Epidemiology*, 31, 125–136.
- Ludvigsson, J. F., Andersson, E., Ekbom, A., Feychting, M., Kim, J.-L., Reuterwall, C., ... Olausson, P. O. (2011). External review and validation of the Swedish national inpatient register. *BMC Public Health*, 11, 450.
- Ludvigsson, J. F., Otterblad-Olausson, P., Pettersson, B. U., & Ekbom, A. (2009). The Swedish personal identity number: Possibilities and pitfalls in healthcare and medical research. *European Journal of Epidemiology*, 24, 659–667.
- Mårild, K., Størdal, K., Bulik, C. M., Rewers, M., Ekbom, A., Liu, E., & Ludvigsson, J. F. (2017). Celiac disease and anorexia nervosa: A nationwide study. *Pediatrics*, 139, e20164367.
- McCarthy, M. M., Nugent, B. M., & Lenz, K. M. (2017). Neuroimmunology and neuroepigenetics in the establishment of sex differences in the brain. *Nature Reviews Neuroscience*.
- Mechawar, N., & Savitz, J. (2016). Neuropathology of mood disorders: Do we see the stigmata of inflammation? *Translational Psychiatry*, 6, e946.

- Monteleone, P., Maes, M., Fabrazzo, M., Tortorella, A., Lin, A., Bosmans, E., ... Maj, M. (1999). Immunoendocrine findings in patients with eating disorders. *Neuropsychobiology*, 40, 115–120.
- Muskens, J. B., Velders, F. P., & Staal, W. G. (2017). Medical comorbidities in children and adolescents with autism spectrum disorders and attention deficit hyperactivity disorders: A systematic review. *European Child & Adolescent Psychiatry*, 26, 1093–1103.
- Mustelin, L., Hedman, A. M., Thornton, L. M., Kuja-Halkola, R., Keski-Rahkonen, A., Cantor-Graae, E., ... Bulik, C. M. (2017). Risk of eating disorders in immigrant populations. *Acta Psychiatrica Scandinavica*, 136, 156–165.
- Neigh, G. N., & Ali, F. F. (2016). Co-morbidity of PTSD and immune system dysfunction: Opportunities for treatment. *Current Opinion in Pharmacology*, 29, 104–110.
- Nielsen, P. R., Benros, M. E., & Dalsgaard, S. (2017). Associations between autoimmune diseases and attention-deficit/hyperactivity disorder: A nationwide study. *Journal of the American Academy of Child and Adolescent Psychiatry*, 56, 234–240.
- Nikolajczyk, B. S., Jagannathan-Bogdan, M., Shin, H., & Gyurko, R. (2011). State of the union between metabolism and the immune system in type 2 diabetes. *Genes and Immunity*, 12, 239–250.
- Okada, Y., Wu, D., Trynka, G., Raj, T., Terao, C., Ikari, K., ... Plenge, R. M. (2014). Genetics of rheumatoid arthritis contributes to biology and drug discovery. *Nature*, 506, 376–381.
- Postler, T. S., & Ghosh, S. (2017). Understanding the holobiont: How microbial metabolites affect human health and shape the immune system. *Cell Metabolism*, 26, 110–130.
- Raevuori, A., Haukka, J., Vaarala, O., Suvisaari, J. M., Gissler, M., Grainger, M., ... Suokas, J. T. (2014). The increased risk for autoimmune diseases in patients with eating disorders. *PloS One*, 9, e104845.
- Rooks, M. G., & Garrett, W. S. (2016). Gut microbiota, metabolites and host immunity. *Nature Reviews Immunology*, 16, 341–352.
- Saucillo, D. C., Gerriets, V. A., Sheng, J., Rathmell, J. C., & Maciver, N. J. (2014). Leptin metabolically licenses T cells for activation to link nutrition and immunity. *Journal of Immunology*, 192, 136–144.
- Smedby, B. (2006). *Health classifications in the Nordic countries: Historic development in a national and international perspective*. Nordisk Medicinalstatistisk Komité.

- Solmi, M., Veronese, N., Favaro, A., Santonastaso, P., Manzato, E., Sergi, G., & Correll, C. U. (2015). Inflammatory cytokines and anorexia nervosa: A meta-analysis of cross-sectional and longitudinal studies. *Psychoneuroendocrinology*, *51*, 237–252.
- Stice, E., Marti, C. N., & Rohde, P. (2013). Prevalence, incidence, impairment, and course of the proposed DSM-5 eating disorder diagnoses in an 8-year prospective community study of young women. *Journal of Abnormal Psychology*, *122*, 445–457.
- Straub, R. H., & Cutolo, M. (2016). Glucocorticoids and chronic inflammation. *Rheumatology*, *55*, ii6–ii14.
- Statistics Sweden. (2012). Longitudinal integration database for health insurance and labour market studies (LISA by Swedish acronym).
- Symmons, D. P. M. (2002). Epidemiology of rheumatoid arthritis: Determinants of onset, persistence and outcome. *Clinical Rheumatology*, *16*, 707–722.
- Theofilopoulos, A. N., Kono, D. H., & Baccala, R. (2017). The multiple pathways to autoimmunity. *Nature Immunology*, *18*, 716–724.
- Tokatly Latzer, I., Rachmiel, M., Zuckerman Levin, N., Mazor-Aronovitch, K., Landau, Z., Ben-David, R. F., ... Pinhas-Hamiel, O. (2018). Increased prevalence of disordered eating in the dual diagnosis of type 1 diabetes mellitus and celiac disease. *Pediatric Diabetes*.
- Tylee, D. S., Sun, J., Hess, J. L., Tahir, M. A., Sharma, E., Malik, R., ... Glatt, S. J. (2017, August 13). *Genetic correlations among neuro-behavioral and immune-related phenotypes based on genome-wide association data. bioRxiv.*
- Ungaro, R., Mehandru, S., Allen, P. B., Peyrin-Biroulet, L., & Colombel, J.-F. (2017). Ulcerative colitis. *Lancet*, *389*, 1756–1770.
- Volk, D. W. (2017). Role of microglia disturbances and immune-related marker abnormalities in cortical circuitry dysfunction in schizophrenia. *Neurobiology of Disease*, *99*, 58–65.
- Wotton, C. J., James, A., & Goldacre, M. J. (2016). Coexistence of eating disorders and autoimmune diseases: Record linkage cohort study, UK. *International Journal of Eating Disorders*, *49*, 663–672.

- Yao, S., Kuja-Halkola, R., Thornton, L. M., Runfola, C. D., D'Onofrio, B. M., Almqvist, C., ... Bulik, C. M. (2016). Familial liability for eating disorders and suicide attempts: Evidence from a population registry in Sweden. *JAMA Psychiatry*, 73, 284–291.
- Yilmaz, Z., Hardaway, J. A., & Bulik, C. M. (2015). Genetics and epigenetics of eating disorders. *Advances in Genomics and Genetics*, 5, 131–150.
- Zallot, C., Quilliot, D., Chevaux, J.-B., Peyrin-Biroulet, C., Guéant-Rodriguez, R. M., Freling, E., ... Peyrin-Biroulet, L. (2013). Dietary beliefs and behavior among inflammatory bowel disease patients. *Inflammatory Bowel Diseases*, 19, 66–72.
- Zerwas, S., Larsen, J. T., Petersen, L., Thornton, L. M., Quaranta, M., Koch, S. V., ... Bulik, C. M. (2017). Eating disorders, autoimmune, and autoinflammatory Disease. *Pediatrics*, 140.